

LexaDerm®EQ

Topical Ointment

Specifications

DESCRIPTION: LexaDerm®EQ serves as an enzymatic debridement agent to slough off any necrotic tissue. It also has a moisturizing agent that keeps the wound bed moist. LexaDerm®EQ opens the surrounding capillaries which helps draw blood flow to the wound site. LexaDerm®EQ demonstrates effective anti-bacterial, anti-fungal and anti-inflammatory responses as well as rapid skin regeneration promotion. LexaDerm®EQ promotes granulation tissue and re-epithelialization of the wound.

CLINICAL PHARMACOLOGY: LexaDerm®EQ demonstrates an effective anti-bacterial and anti-inflammatory response as well as rapid skin regeneration promotion.

INDICATIONS AND USAGE: An external treatment for the inflammation and irritation associated with many common forms of dermatitis, including certain eczematoid conditions. These conditions include complications associated with pyodermas. Also indicated in the treatment of insect bites, burns and fungal infections.

CONTRAINDICATIONS: No known.

WARNINGS: A mild itching or warming sensation may be experienced during treatment. For external use only. Avoid contact with eyes. If irritation occurs or symptoms persist after 10 days, discontinue use and consult your veterinarian.

OTHER INFORMATION:

Patients should be advised that:

1. Lexaderm®EQ should be applied daily or every 12 hours.
2. It is for external use only. Avoid contact with the eyes.

DRUG INTERACTIONS

It is not known if LexaDerm®EQ interacts with other topical medications applied to the ulcer site. The use of LexaDerm®EQ with other topical drugs has not been studied.

PREGNANCY

Pregnancy studies in humans and animals have not been done to determine whether LexaDerm®Eq is teratogenic. However, there is no evidence that moderate therapeutic doses of Lexaderm®EQ causes fetal damage in humans.

DOSAGE AND ADMINISTRATION:

For moderate skin conditions: Wash area to be treated, pat dry, apply LexaDerm®EQ to the area to be treated once daily or as directed by veterinarian. Keep area dry. To protect clothing, area may be bandaged. Use light dressing or gauze. Follow good wound care procedures with each treatment.

For open wounds: Remove old dressing and discard. Flush wound with 250-500 cc normal saline (if patient unable to shower). If patient is unable to shower, instruct to flush wound with tepid water for 5-10 minutes. Apply a generous amount of : LexaDerm®EQ with a cotton-tipped applicator or other application aid to wound base only. Apply skin protectant to surrounding skin. Fluff dressing and apply to dry wound. Cover with transparent dressing. Change daily or q 12 hours.

HOW SUPPLIED: LexaDerm®EQ topical is supplied in the following sizes:

- 1 oz. Tube
- 4 oz. Tube

STORAGE:

Store at room temperature, 59-86 degrees Fahrenheit.

Reference: 1. Data on file, 7 Oaks Pharmaceutical Corporation.

Composition of : LexaDerm®EQ

Actives: Benzoic Acid, USP - 6.0%

With: Polyethylene Glycol-400

Polyethylene Glycol – 3350

QRB7 – 3% (extract of oak bark)